

VII CONGRESO URUGUAYO DE OBESIDAD

1º Congreso Regional de Obesidad y Trastornos Alimentarios

**Balance Energético y Regulación
Siconeuroendócrina de la Ingesta**

Dr Fernando M. Escobar
Médico Endocrinólogo Universitario
Ex Presidente de la Sociedad Argentina
de Obesidad y Trastornos Alimentarios
Docente Adscripto en Medicina Interna

- El relator de la conferencia no presenta conflictos de interés.



Daniel Lambert

1770-1809

335 kg

"The most corpulent man of whom authentic records exist"

Died age 39, Heart Disease






**“A Voluptuary under the
Horrors of Digestion”**

1792, James Gillray

*Prince Regent
(Future King George IV of
England)*

Etiología de la Obesidad



**Ingreso
calórico**

**Gasto
Calórico**

Sedentarismo

**Alimentos de alta
densidad calórica
azúcar y grasas**

**Susceptibilidad
Genética**

Mecanismos del ingreso alimentario

Influencias familiares y sociales



Factores sensoriales

- Sabor
- Olor
- Textura
- Efecto visual
- **Efectos de**
- Variedad
- Saciedad al estímulo
- Sensorial
- Palatabilidad
- Tipo de alimento
- Disponibilidad real



Factores cognitivos

- Control conciente
- Creencias alimentarias
- Advertencias



Mecanismos cerebrales

- Modula factores sensoriales mediante señales de saciedad para producir sensación de recompensa y regulación del apetito.




Señales de Saciedad y Apetito


- Hormonas del tej. Adiposo
- Hormonas intestinales
- Distensión gástrica

Ingesta

Los diferentes niveles de control del apetito




Neuroregulación del balance energético





Amylin



GLP-1



Oxyntomodulin




PYY₃₋₃₆




Courtesy Hans-Rüdiger Barthold




Leptin



Péptidos del tubo digestivo que regulan el apetito



GLP-1



- Péptido de 30 aa producido a partir del pre-proglucagon en el NTS y cel.L Intestinales.
- GLP-1 se produce e incluso activa neuronas del NTS que se proyectan a regiones claves del hipotálamo.
- GLP-1 disminuye la ingesta en ratas. El antagonista GLP-1R Exendina aumenta el apetito.
- La administración de GLP-1 activa neuronas en el núcleo arcuato, PVN, NTS, y área postrema.
- Inhibe el apetito

Mecanismos potenciales para GLP.1 que median en la disminución de la ingesta y del peso corporal

- **Enlentecimiento del vaciado gástrico**

- **Saciedad**

 - ***Directo*** (Vía NTS y/o hipotálamo)

 - ***Indirecto*** (Vía Aferentes vagales)




Otras acciones del GLP-1



- **Segregado por las células L-intestinales con la ingesta**
- **En humanos y en animales**
 - **Incrementa la liberación de insulina estimulada por glucosa**
 - **Disminuye la secreción de glucagon**
 - **Disminuye el vaciado gástrico**
 - **Reduce el apetito**
- **En animales in vitro**
 - **Aumenta la transcripción del gen de insulina**
 - **Aumenta la masa y diferenciación de las células β**

Drucker DJ. *Curr Pharm Des.* 2001;7:1399-1412.
Drucker DJ. *Mol Endocrinol.* 2003;17:161-171.

GLP-1 Reduce la ingesta alimentaria





* $P < 0.05$, † $P < 0.001$, vs control.
Gutzwiller JP, et al. Gut 1999;44:51-56.

Efecto de un GLP-1 de acción prolongada sobre glucosa y balance energético

HbA1c

Peso corporal



Glicentina y Oxyntomodulina

Glicentina

- -Hormona intestinal producida por las células α y L
- -Liberada durante la digestión
- -Insulino-trópica, disminuye la acidez y el vaciado gástrico
- -No hay todavía data disponible en humanos

Oxyntomodulina

- -37 aa, igual al glucagon más su extensión C terminal
- -Segregado por las células L, luego de la ingesta, se une al receptor GLP-1
- -Reduce la ingesta alimenticia en humanos
- -Puede aumentar el gasto energético ¿vía el eje tiroideo?


Acciones centrales de PYY₃₋₃₆ y Oxyntomodulina

OXM y PYY₃₋₃₆ se liberan postprandialmente


-Acción primaria: señales de saciedad en el núcleo arcuato.

-Inhibición coordinada de neuronas orexígenas y activación de anorexígenas

-Reducción del gasto energético y del peso corporal.



Impacto de la Oxintomodulina en la ingesta alimentaria y en el peso corporal




Cohen et al. JCEM 38(10):4696-4703, 2003
Wynne et al. Diabetes 54, August 2005
Wynne et al. Int J Obesity 2006 30, 1729-1736

13.06.2009 15:16

Las hormonas leptina e insulina ingresan al núcleo arcuato hipotalámico para estimular las neuronas α -MSH/CART mientras que inhiben NPY/AgRP.

Por lo tanto a nivel central activan los sistemas catabólicos e inhiben los anabólicos.



Ghrelin: es un ligando endógeno del receptor secretagogo de GH


Se expresa en estómago

- Estimula la secreción de GH

- Incrementa la adiposidad disminuyendo la oxidación de las grasas y aumentando la ingesta

- Los niveles de Ghrelin están disminuídos en obesos

- Ghrelin disminuye luego de una cirugía de bypass gástrico




Adapted from Cummings et al. NEJM 346:1623-1630, 2002

Receptor MC4R e ingresso alimentare

Leptin $\uparrow$ POMC
and $\downarrow$ AGRP
 $\rightarrow \uparrow$ MC4R
signaling and $\downarrow$
food intake


MC4R signaling
is $\uparrow$ by α -MSH
and $\downarrow$ by AGRP




Leptin $\downarrow$ NPY, which is synthesized in the same neurons as AGRP

Stimulatory interactions in solid lines; inhibitory in dashed lines
Adapted from SL Wardlaw JCEM 86:1442-1446, 2001

Los Endocannabinoides estimulan los circuitos anabólicos e Inhiben los circuitos catabólicos en el hipotálamo




Endocannabinoids Increase Food Intake Through CB₁-Receptor Activation




Anandamide administration into the ventromedial hypothalamus induces hyperphagia in presatiated rats

Leptin tonically inhibits endocannabinoids in the hypothalamus.




Effects of Endocannabinoid System Overactivity



¿Cuál es la Causa de la Epidemia de Obesidad?

La comida "Basura" provoca Leptino-resistencia



Adapted from Campfield LA, et al. Science. 1998;280:1383-1387; Porte D, et al. Diabetologia. 1998;41:863-881.

Endoplasmic Reticulum Stress Plays a Central Role in Development of Leptin Resistance

Lale Ozcan,^{1,*} Ayse Seda Ergen,^{1,*} Allen Lu,² Jason O'Hong,² Sumit Sarkar,² Danyu Niu,² Martin G. Myers, Jr.,² and Umut Ozcan^{1,2}

¹Division of Endocrinology

²Division of Neurology

Children's Hospital Boston, Harvard Medical School, Boston, MA, 02115, USA

²Division of Metabolism, Endocrinology, and Diabetes, Department of Medicine, University of Michigan Medical School, Ann Arbor, MI, 48106, USA

*These authors contributed equally to this work.

Correspondence: umut.ozcan@childrens.harvard.edu

DOI: 10.1016/j.cmet.2009.12.004

- **La alimentación engordante causa Leptino-resistencia**
- **El incremento de peso provoca Leptino-resistencia**
- **La Leptino- resistencia interfiere con la saciedad y altera el metabolismo del tejido muscular favoreciendo el incremento de peso**
- **Los malos hábitos generan también una “Mala Fsiología”**

Endoplasmic Reticulum Stress Plays a Central Role in Development of Leptin Resistance

Lale Ozcan,^{1,*} Ayse Seda Ergen,^{1,*} Allen Lu,² Jason Chung,² Sumit Sarkar,² Dany Niu,² Martin G. Myers, Jr.,² and Umut Ozcan^{1,2}

¹Division of Endocrinology

²Division of Neurology

Children's Hospital Boston, Harvard Medical School, Boston, MA 02115, USA

²Division of Metabolism, Endocrinology, and Diabetes, Department of Medicine, University of Michigan Medical School, Ann Arbor, MI 48106, USA

*These authors contributed equally to this work.

Correspondence: umut.ozcan@childrens.harvard.edu

DOI: 10.1016/j.cmet.2009.12.004


- “Taking all of these phenomenons into account, we think that it is possible that overconsumption of nutrients could itself be a reason for development of leptin resistance”
- “This line of thinking favors the fact that increased adiposity and consequent hyperleptinemia decreases the leptin action and creates the leptin resistance”

Otras Areas del Cerebro están Involucradas además del Hipotálamo

Los Circuitos de Recompensa también Influyen en la Conducta Alimentaria

**Sistema mesolímbico
Dopaminérgico**

**Adicción
Placer (opioides)
Deseo (dopamina)**



**Ingesta por palatabilidad
Motivación**

Dopamina (bupropion: tratamiento para dejar de fumar)

Opioides (Naltrexona: utilizada para tratar la dependencia opioide y alcohólica)

Neurobiología del apetito apetito como adicción

- **Amígdala, insula, corteza orbito-prefrontal y striatum son elementos Centrales en el control de la conducta alimentaria.**
 - ✓ **Aprendizaje en relación a la recompensa alimentaria**
 - ✓ **Prestar atención y esfuerzo para lograr dicha recompensa**
 - ✓ **Cuál es el valor que se le otorga ese estímulo en el medio ambiente**
 - ✓ **Integrar esa información en relación a la reserva energética y a la disponibilidad alimentaria**




Figure 2. Activation in a region of the anterior cingulate cortex (Montreal Neurological Institute coordinates x, y, z : $-9, 24, 27$; $z=4.64$; false discovery rate-corrected $P.001$) during milkshake cues vs tasteless solution cues as a function of Yale Food Addiction Scale scores, with the graph of parameter estimates (PE) from that peak.





Figure 3. Activation in a region of the medial orbitofrontal cortex (Montreal Neurological Institute coordinates x, y, z : 3, 42, -15; $z=3.47$; false discovery rate-corrected $P=.004$) during milkshake cues vs tasteless solution cues as a function of Yale Food Addiction Scale scores, with the graph of parameter estimates (PE) from that peak




Figure 4. Activation in a region of the dorsolateral prefrontal cortex (DLPFC) (Montreal Neurological Institute coordinates x, y, z : $-27, 27, 36$; $z=3.72$; false discovery rate–corrected $P=.007$) during anticipatory food reward (milkshake cue vs tasteless solution cue) in the high food addiction (FA) group vs the low FA group, with the bar graphs of parameter estimates (PE) from that peak.




Figure 6. Activation in a region of the lateral orbitofrontal cortex (Montreal Neurological Institute coordinates x, y, z : $-42, 42, -12$; $z = -3.45$; false discovery rate-corrected $P = .009$) during consummatory reward (milkshake receipt vs tasteless solution receipt) in the high food addiction (FA) group vs the low FA group, with the bar graphs of parameter estimates (PE) from that peak.

- La extirpación de la amígdala anula la atracción por las características del alimento ej. Aroma
- Dopamina proveniente del cerebro mesolímbico produce efecto hedónico
- Ghrelina refuerza mientras que insulina y leptina inhiben el deseo

- **Algunos individuos reaccionan con más impulsividad ante ciertos alimentos (Caroline Davis)**
- **Los opioides promueven los alimentos de alta densidad energética estimulan el deseo y lentifican la saciedad**
- **La amígdala recibe múltiples estímulos tales como textura, aromas, sabores, temperatura y controles de las hormonas peptídicas intestinales.(su lesión quita estímulo de los alimentos muy calóricos y las ganas de fumar)**

- **La lesión del área prefrontal o su atrofia produce hiperfagia a veces con preferencia por alimentos más elaborados (síndrome del gourmand)**
- **La hiperactividad (epilepsia) provoca un símil anorexia nerviosa que cede con la terapia anticonvulsivante**
- **El área prefrontal derecha elige mejor los alimentos cuanto mayor es su desarrollo. Los obesos desarrollan más el hemisferio izquierdo,**
- **También influye en la percepción de la imagen corporal.**
- **La disfunción del area prefrontal derecha por atrofia provoca incremento de la alimentación, sobretodo fast food**


Modelo cerebral de la obesidad humana

- **La alimentación en los humanos se puede dividir en refleja o reflexiva**
- **La refleja depende de los circuitos neurohipotalámicos y de los neuropéptidos que controlan el apetito y la saciedad.**
- **Por encima se ubica la reflexiva que involucra conductas relacionadas con criterios sociales y culturales.**
Cultura del alimento sano, de la figura esbelta, etc.

Mecanismo hipotético de “Alimentación Progresiva” para generar un continuo Incremento de peso



El futuro: Tratamientos combinados





**¡Gracias por
haberme
escuchado!**



Liraglutide Effects on Glycemic Control and Body Weight

Liraglutide (qd) ■ 0.045 mg ○ 0.225 mg ▲ 0.45 mg ■ 0.60 mg ● 0.75 mg

Metformin ◆ 1,000 mg bid




*Mean baseline A1C for all groups: 6.8% to 7.5%.

†Significant difference compared to metformin.

Feinglos MN et al. *Diabet Med.* 2005;22:1016-1023.


HYPOTHALAMUS






LEAD-4: Effect of Liraglutide in Combination With Metformin and Rosiglitazone on Body Weight in Patients With Type 2 Diabetes

- Liraglutide (1.2 mg/qd) + metformin (1 g/bid) + rosiglitazone (4 mg/bid) (n=178)
- Liraglutide (1.8 mg/qd) + metformin (1 g/bid) + rosiglitazone (4 mg/bid) (n=178)
- Placebo + metformin (1 g/bid) + rosiglitazone (4 mg/bid) (n=177)



^a $P < 0.05$ vs placebo.

Zinman B et al. *Diabetologia*.
2008;51(suppl 1):Poster 898.







Glucagon-Like Peptide-1



- Secreted from intestinal L-cells with meal ingestion
- In humans and animals
 - enhances glucose-stimulated insulin release
 - decreases glucagon release
 - slows gastric emptying
 - reduces food intake
- In animals and in vitro
 - increases insulin gene transcription
 - increases β -cell mass and β -cell differentiation




Once-Weekly vs Twice-Daily Exenatide in Type 2 Diabetes: Body Weight




* Exenatide's once-weekly dosage is investigational. LAR=long-acting release.

Drucker DJ et al. [Published online ahead of print September 8, 2008]. *Lancet*. doi:10.1016/S0140-6736(08)61206-4.



Change in Body Weight Following 82 Weeks of Exenatide Treatment




Blonde L et al. *Diabetes Obes Metab.* 2006;8:436-447.

Liraglutide

- -Una dosis diaria es efectiva
- -Posee efectos beneficiosos en la glucemia y peso corporal en pacientes
- diabéticos
- -Estimula la liberación de insulina, suprime la gluconeogénesis y provoca
- náuseas-

Liraglutide altered food preference: decreased chocolate but increased chow intake




Raun, Vess, Gottfredsen, Golozoubova, Rolin & Bjerre Knudsen, Diabetes 2007;56:8-15


13.06.2009 20:55

Complementary Central Action in Obesity PYY₃₋₃₆ and Oxyntomodulin


- OXM and PYY₃₋₃₆ released post-prandially
 - Primary action → as satiety signals in hypothalamic arcuate nucleus
 - Coordinated inhibition of *orexigenic* neurons and activation of *anorectic* neurons
- Animal/human studies
 - Reduced energy intake and weight loss




Leptin/Endocannabinoid Interaction in the Lateral Hypothalamus



Hyperphagia from loss of
leptin signaling




Normalization by CB₁R
antagonists (rimonabant)



Reproduced from Jo YH, et al. *Neuron*.
2005;48:1055-1066.

The control system over caloric homeostasis is complex, relying upon several types of signals



Schwartz MW, et al. *Nature*. 2000;404:661-671.

How does the body control its own weight?

1 ton of food = 1 million calories in 1 year




Genetics and Environment

Sweet, high fat foods
Hormones to suppress appetite
Storing fat; energy expenditure

Alcohol; smoking
Excess calories
Physical inactivity

Obesity Treatment Pyramid



Multi-disciplinary Approach to Obesity Management


Weight Management Care Team




Optimal Weight Loss Goals

- Optimal rate of weight loss is 1-2 kg, or 2-4 pounds, a month.
- Aim for a deficit of 500 kcal/day to achieve a weight loss of a pound (or 3500 kcal) a week
- A 500 kcal deficit diet would amount to approximately 20-30% decrease in average daily food intake


What is Success in Weight Management?



Age-related Weight Gain Cannot Be Prevented Despite Continued Intervention Efficacy



No quick fixes!




"Just pull my sweet tooth"

Source: 1989 H.L. Schwadron

Novel Anti-obesity Drugs

- Appetite suppressants that target
 - Leptin (recombinant leptin)
 - Neuropeptide Y
 - Melanocortin-4 receptor
 - Cannabinoid receptors (CB1 antagonist-rimonabant)
 - Dopamine (ecopipam)
- Ghrelin antagonist
- PYY3-36
- Ciliary neurotrophic factor – Axokine
- Selective β_3 adrenergic receptor agonists to increase energy expenditure
- Fat mobilization via lipolysis, inhibitors of fat digestion and absorption
- Nutrient partitioning to favour lean body mass



Is leptin the brain signal?


Leptin is a fat-derived lipostat signaling hormone

Recombinant Leptin Therapy


- Randomized, double-blind, placebo-controlled trial in 54 lean and 73 obese adults (BMI 30-36) for 4-24 weeks
 - Dose-dependent weight loss in obese subjects
 - 0.7 to 7.1 kg
- Antileptin antibodies present in 38% and 71% at 4 and 24 weeks
- Injection site reactions common
- Majority of obese subjects are leptin resistant!

Adapted from Heymsfield et al, JAMA 282:1568-1575, 1999


Regulation of Appetite



Cannabinoid Receptors




(-)- Δ^9 -tetrahydrocannabinol
(THC, oil)



2-Arachidonoylglycerol



- Hippocampus
- Basal Ganglia
- Cortex
- Cerebellum
- Hypothalamus
- Limbic structures
- Brainstem
- GI tract
- Immune cells & tissues



The Endocannabinoid System is a Modulatory System


■ Endocannabinoids:

- Synthesized on demand from lipid precursors in postsynaptic cell
- Activate CB_1 receptors presynaptically, then degraded immediately
- Act as retrograde messengers
- Inhibit neurotransmitter release


■ CB_1 receptors:

- Play a key role in food intake, energy balance and lipid and glucose metabolism


V Di Marzo 2005
V Di Marzo 1998
R Wilson 2002



Rimonabant (CB₁ Blocker)




Central and Peripheral Targets of ECS and Effects of Overactivity




M Bensaid M 2003; U Pagotto 2005; D Osei-Hyiaman 2005; V DiMarzo 2005; YL Liu 2005

The only way to keep healthy is to eat what you don't want, drink what you don't like and do what you would rather not do

Mark Twain (1835 - 1910)





Leptin




Oxyntomodulin

PYY₃₋₃₆

GLP-1



Amylin



GLP-1 Processing




Prohormone Convertase-1


Inactive GLP-1

DPP-4

GLP-1 and -2=Glucagon-like peptide-1 and -2; GRPP=Glucagon-related pancreatic peptide; IP-1 and -2=Interleukin peptides 1 and 2.
Redrawn from Moller DE. *Nature*. 2001;414:821-827


● GLP-1 Receptors





Ma X, et al, J Neurosci. 2007 Jul 4;27(27):7125-9.

Placebo-Corrected Change in Weight





Data summarized from results reported in abstracts from the 68th sessions of the American Diabetes Association, 2008.

Liraglutide and exenatide have not been studied in head to head trials.

*Changes in weight for trials with no placebo arm are shown as change from baseline.


Weight Reductions With 3.5 Years of Exenatide



3.5-year completer cohort (n=151). Baseline weight 90.9 kg. Mean ± SE.
Kendall D, et al. Diabetes. 2007;56(suppl 1):A149. Data on file, Amylin Pharmaceuticals, Inc.













Adiposity Signals: Leptin




Neuroregulation of Energy Balance






Regulation of Appetite







Gut Peptides That Regulate Appetite



COR-I, II: Body Weight,

Percentage of Change From




NB = naltrexone/bupropion. #COR-II: NB observed data are NB32/NB48 pooled (N = 825); no differences were observed for patients rerandomized to NB32 vs NB48. LS mean $\pm$ SE; * P < .001 vs placebo at all timepoints. COR-II: week 56 ITT-LOCF data from patients rerandomized to NB32 are double weighted to account for the prespecified exclusion of patients rerandomized to NB48. ITT-LOCF: patients with a baseline and ≥ 1 postbaseline weight measurement while on study drug.

Press release. October 27, 2009. Available at:

<http://ir.orexigen.com/phoenix.zhtml?c=207034&p=irol-newsArticle&ID=1346886&highlight> Accessed April 29, 2010.



Modelo simplificado de los mecanismos de regulación del peso corporal



Circuitos metabólicos periféricos del balance energético

Las señales para comer o no comer que se dirigen al hipotálamo provienen de:

- **Tubo digestivo**
- **Páncreas**
- **Tejido Adiposo**

